arXiv:1403.6999v2 [hep-th] 31 Mar 2014

A class of hermitian generalized Jordan triple systems and Chern-Simons gauge theory

Noriaki Kamiya *
Department of Mathematics, University of Aizu
Aizuwakamatsu, 965-8580, Japan
Matsuo Sato †
Department of Natural Science, Faculty of Education, Hirosaki University
Bunkyo-cho 1, Hirosaki, Aomori 036-8560, Japan

Abstract

We find a class of hermitian generalized Jordan triple systems (HGJTSs) and hermitian (ϵ, δ)-Freudenthal-Kantor triple systems (HFKTSs). We apply one of the most simple HGJTSs which we find to a field theory, and obtain a typical $u(N)$ Chern-Simons gauge theory with a fundamental matter.

AMS classification: 17C50; 17A40; 17B60 Keywords: generalized Jordan triple systems, Lie algebras, hermitian operators.
APS classification: 11.15.Yc; 11.25.Yb; 11.25.-w Keywords: 3-algebras, Chern-Simons gauge theories, M-theory

Introduction

HGJTSs and HFKTSs [1, 2] are generalizations[1] of hermitian 3-algebras [6–50], which have played crucial roles in M-theory. The field theories applied with hermitian 3-algebras are the Chern-Simons gauge theories with bi-fundamental matters that describe the effective actions of membranes in M-theory[2]. Moreover, 3-algebra models of M-theory itself have been proposed and were studied in [53–60]. Therefore, it is interesting to apply HGJTSs and HFKTSs to field theories.

In [1, 2], we searched HGJTSs and HFKTSs by adding some terms to the Hermitian 3-algebras. In this paper, we study a class of HGJTSs and HFKTSs systematically. We apply one of the most simple HGJTSs which we find to a field theory, and obtain a typical $u(N)$ Chern-Simons gauge theory with a fundamental matter.

Here, we are concerned with algebras and triple systems that are finite or infinite over a complex number field, unless otherwise specified. This paper is an announcement of

*e-mail address : kamiya@u-aizu.ac.jp
†e-mail address : msato@cc.hirosaki-u.ac.jp

[1] A different generalization, the so-called differential crossed module, was extensively investigated in [3–5].

[2] The structures of generalized Jordan triple systems were found in the effective actions of the membranes in [51, 52]

new results, that is first step of our works for hermitian triple systems and applications to field theories. hence we will give a lot of references [61–75]. We refer to [76–80] for nonassociative algebras, and to [81–83] for Lie superalgebras, for example.

1 Definitions

In this section, we summarize definitions of HGJTS and HFKTS given in [1, 2].

A triple system U is a vector space over a field F of characteristic $\neq 2, 3$ with a trilinear map $U \otimes U \otimes U \to U$. In this paper, we are concerned with triple systems over the complex number $\mathbf{C}$, and denote the trilinear product by $< xyz > \in U$ for $x, y, z \in U$, assuming that $< xyz >$ is $\mathbf{C}$-linear on x and z, and $\mathbf{C}$-antilinear on y. Also we use the notations of two operations $L(x, y) \in \text{End } U$ and $K(x, y) \in \text{End } U$ with respect to the triple systems ([1], [2]), where $L(x, y)z =< xyz >$ and $K(x, y)z =< xzy > -\delta < yzx > \quad (\delta = \pm 1)$.

Definition. A triple system U is said to be a *-(ϵ, δ)-Freudenthal-Kantor triple system (FKTS) if relations (0)–(iv) are satisfied.

0) U is a Banach space,

i) $[L(x, y), L(z, w)] = L(< xyz >, w) + \varepsilon L(z, < yxw >)$ (fundamental identity), where $[A, B] = AB - BA$, $A, B \in \text{End } U, and \ \varepsilon = \pm 1$,

ii) $K(< xyz >, w) + K(z, < xyw >) + \delta K(x, K(z, w)y) = 0$ (second order identity),

iii) $< xyz >$ is a $\mathbf{C}$-linear operator on x and z, and a $\mathbf{C}$-antilinear operator on y,

iv) $< xyz >$ is continuous with respect to a norm, $|| \ ||$ that is, there exists $K > 0$ such that

$$|| < xxx > || \leq K||x||^3 \text{ for all } x \in U.$$

U is called *-generalized Jordan triple system (GJTS) if 0), i), iii), and iv) are satisfied when $\epsilon = -1$ and $\delta = -1$. The condition ii) is called second order identity and *-generalized Jordan triple system of second order is (-1, -1) FKTS. Note that there are many simple Lie algebras (the case of $\delta = 1$) and simple Lie superalgebras (the case of $\delta = -1$) constructed from (ϵ, δ)-Freudenthal-Kantor triple systems ([4,7,8]) because of the second order identity.

Furthermore, a *-(ϵ, δ)-FKTS and *-GJTS are said to be hermitian if they satisfy the following condition.

v) All operators $L(x, y)$ are operators with a hermitian metric

$$(x, y) = tr \ L(x, y),$$

that is, $(L(x, y)z, w) = (z, L(y, x)w)$ and $(x, y) = \overline{(y, x)}$.

2 A class of HGJTSs and HFKTSs

In this section, we obtain a class of HGJTSs and HFKTSs by making an ansatz $< xyz >= \alpha x\tilde{y}z + \beta z\tilde{y}x + \gamma \bar{y}\tilde{\tilde{x}}z$ and explicitly evaluating the conditions i) and ii) in the last section. The hermitian 3-algebra $x\tilde{y}z - z\tilde{y}x$ is in a special case when $\alpha = 1$, $\beta = -1$, and $\gamma = 0$. By straightforward, but extensive calculations (to be omitted the proof here), we obtain theorems as follows.

Theorem A. Let U be a hermitian triple system over complex number $\mathbf{C}$ satisfying the triple product $< xyz >= \alpha x\tilde{y}z+\beta z\tilde{y}x+\gamma\bar{y}\tilde{\tilde{x}}z$ for $\alpha, \beta, \gamma \in \mathbf{R}$ (real number), $x, y, z \in U$, induced from an associative algebra with an involution $\widetilde{xy} = \tilde{y}\tilde{x}$ and $\tilde{\tilde{x}} = x$.

Then we have a necessary and sufficient condition for U to be a hermitian generalized Jordan triple system is that the following hold;

$\gamma(\alpha + \gamma) = 0$.

Furthermore, we have for U to have the second order condition ii) is the following holds;

$\alpha = \beta$ if $\delta = 1$,

$\alpha = -\beta$ if $\delta = -1$.

Summarizing, we obtain a class of HGJTSs,

Proposition B. Let U be a set of $M(n, n; \mathbf{C})$ with the triple product defined by

1)$< xyz >$ is a linear combination of $x\tilde{y}z$ and $z\tilde{y}x$,

2) $< xyz >$ is a linear combination of $x\tilde{y}z - \bar{y}\tilde{\tilde{x}}z$ and $z\tilde{y}x$,

where $\tilde{x} = \bar{x}^T, ^-$is the conjugation of $\mathbf{C}$, and x^T is the transpose of x. Then U are HGJTSs. The hermitian 3-algebra $< xyz >= x\tilde{y}z - z\tilde{y}x$ is in a special case of 1). One of the most simple cases, $< xyz >= x\tilde{y}z$ will be applied to a field theory in the next section.

We also obtain a class of HFKTSs,

Proposition C. Let U be a set of $M(n, n; \mathbf{C})$ with the triple product defined by

a) $< xyz >= x\tilde{y}z + z\tilde{y}x$,

b) $< xyz >= x\tilde{y}z + z\tilde{y}x - \bar{y}\tilde{\tilde{x}}z$,

c) $< xyz >= x\tilde{y}z - z\tilde{y}x$,

d) $< xyz >= x\tilde{y}z - z\tilde{y}x + \bar{y}\tilde{\tilde{x}}z$,

where $\tilde{x} = \bar{x}^T, ^-$is the conjugation of $\mathbf{C}$, and x^T is the transpose of x. Then we have the case of a) and b) are (-1,1)-HFKTSs and the case of c) and d) are (-1,-1)-HFKTSs.

Remark. For the case of nonassociative and nonhermitian algebras, we note that there is a structureable algebra due to Allison [76] satisfying the triple product $< xyz >= (x\tilde{y})z + (z\tilde{y})x - (z\tilde{x})y$ and $\widetilde{xy} = \tilde{y}\tilde{x}$, and $\tilde{\tilde{x}} = x$, which is a (-1,1)-Freudenthal-Kantor triple system also this algebra contains a class of commutative Jordan algebras and associative algebras.

We call a hermitian (α, β, γ) structureable algebra about the hermitian triple system induced from a nonassociative algebra satisfying $< xyz >= \alpha(x\tilde{y})z + \beta(z\tilde{y})x + \gamma(z\tilde{\tilde{x}})\bar{y}$. However, the details will be discussed in other paper.

3 Chern-Simons gauge theory with a fundamental matter

In this section, we apply one of the most simple HGJTSs to a field theory. We define a 3-bracket, $[x, \overline{y}, z] :=< xyz >$.

We start with

$$S = \int d^3x \mathrm{tr}(-\mathbf{D}_\mu Z \overline{\mathbf{D}^\mu Z}^T + V(Z) \\ +\epsilon^{\mu\nu\lambda}(-\frac{1}{2}A_{\mu\bar{b}c}\partial_\nu A_{\lambda\bar{d}a}\overline{T}^{T\bar{d}}[T^c, \overline{T}^{\bar{b}}, T^a] \\ +\frac{1}{3}A_{\mu\bar{d}a}A_{\nu\bar{b}c}A_{\lambda\bar{f}e}[T^c, \overline{T}^{\bar{b}}, T^a]\overline{[T^f, \overline{T}^{\bar{e}}, T^d]})), \tag{1}$$

where

$$\mathbf{D}_\mu Z = \partial_\mu Z - A_{\mu\bar{b}a}[T^a, \overline{T}^{\bar{b}}, Z]. \tag{2}$$

Z and A_μ are matter and gauge fields, respectively. $V(Z)$ is a potential term. This action is invariant under the transformations generated by the operator $L(x,y) - L(y,x)$. The action (1) equipped with a Lorentzian Lie 3-algebra and a hermitian 3-algebra describes the bosonic parts of the effective actions of supermembranes in M-theory. We apply a HGJTS, $[x, \overline{y}, z] =< xyz >= x\bar{y}^T z$ in the previous section to this action and see what kind of field theory we obtain.

The covariant derivative is explicitly written down as

$$\mathbf{D}_\mu Z = \partial_\mu Z - iA_\mu Z, \tag{3}$$

where $A_\mu := -iA_{\mu\bar{b}a}T^a\overline{T}^{T\bar{b}}$ are hermitian matrices, which generate the $u(N)$ Lie algebra. In order to see how gauge transformations act on these fields, let us see how $L(x,y) - L(y,x)$ acts on the coupling between A_μ and Z:

$$\begin{aligned} & \delta(A_{\mu\bar{b}a}[T^a, \overline{T}^{\bar{b}}, Z]) \\ = \; & \Lambda_{\bar{d}c}A_{\mu\bar{b}a}([[T^c, \overline{T}^{\bar{d}}, T^a], \overline{T}^{\bar{b}}Z] - [T^a, \overline{[T^d, \overline{T}^{\bar{c}}, T^b]}, Z] + [T^a, \overline{T}^{\bar{b}}, [T^c, \overline{T}^{\bar{d}}, Z]] \\ = \; & [i\Lambda, iA_\mu]Z + iA_\mu i\Lambda Z \\ = \; & i\Lambda iA_\mu Z, \end{aligned} \tag{4}$$

where gauge parameters Λ is defined in the same way as A_μ. From this, we obtain gauge transformations,

$$\begin{aligned} \delta A_\mu &= [i\Lambda, A_\mu] \\ \delta Z &= i\Lambda Z. \end{aligned} \tag{5}$$

Therefore, A_μ transforms as adjoint representations of $u(N)$, whereas Z transforms as a fundamental representation of $u(N)$. As a result, the action can be rewritten in a covariant form and we obtain a typical $u(N)$ Chern-Simons gauge theory with a fundamental matter,

$$\begin{aligned} S &= \int d^3x \mathrm{tr}(-(\partial_\mu Z - iA_\mu Z)\overline{(\partial_\mu Z - iA_\mu Z)}^T + V(Z) \\ & +\epsilon^{\mu\nu\lambda}(\frac{1}{2}A_\mu\partial_\nu A_\lambda + \frac{i}{3}A_\mu A_\nu A_\lambda). \end{aligned} \tag{6}$$

Acknowledgements

The work of M.S. is supported in part by Grant-in-Aid for Young Scientists (B) No. 25800122 from JSPS.

References

[1] N. Kamiya, M. Sato, "Hermitian generalised Jordan triple systems and certain applications to field theory," preprint.

[2] N.Kamiya and M.Sato, "Hermitian (ϵ, δ)-Freudenthal-Kantor triple systems and certain applications of *-generalized Jordan triple systems to field theory," to appear in AHEP

[3] S. Palmer, C. Saemann, "M-brane Models from Non-Abelian Gerbes," JHEP 07 (2012) 010.

[4] S. Palmer, C. Saemann, "Six-Dimensional (1,0) Superconformal Models and Higher Gauge Theory," J. Math. Phys. 54 (2013) 113509.

[5] S. Palmer, C. Saemann, "The ABJM Model is a Higher Gauge Theory," arXiv:1311.1997.

[6] Y. Nambu, Generalized Hamiltonian dynamics, Phys. Rev. D7 (1973) 2405.

[7] V. T. Filippov, n-Lie algebras, Sib. Mat. Zh. 26, No. 6, (1985) 126140.

[8] E. Bergshoeff, E. Sezgin, P.K. Townsend, Supermembranes and Eleven-Dimensional Supergravity, Phys. Lett. B189 (1987) 75.

[9] B. de Wit, J. Hoppe, H. Nicolai, On the Quantum Mechanics of Supermembranes, Nucl. Phys. B305 (1988) 545.

[10] H. Awata, M. Li, D. Minic, T. Yoneya, On the Quantization of Nambu Brackets, JHEP 0102 (2001) 013.

[11] D. Minic, "M-theory and Deformation Quantization," arXiv:hep-th/9909022.

[12] J. Bagger, N. Lambert, "Modeling Multiple M2's," Phys. Rev. D75: 045020, 2007, hep-th/0611108.

[13] A. Gustavsson, "Algebraic structures on parallel M2-branes," Nucl. Phys. B811: 66, 2009, arXiv:0709.1260 [hep-th].

[14] J. Bagger, N. Lambert, "Gauge Symmetry and Supersymmetry of Multiple M2-Branes," Phys. Rev. D77: 065008, 2008, arXiv:0711.0955 [hep-th].

[15] O. Aharony, O. Bergman, D. L. Jafferis, J. Maldacena, "N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals," JHEP 0810: 091, 2008, arXiv:0806.1218 [hep-th]

[16] J. Bagger, N. Lambert, "Three-Algebras and N=6 Chern-Simons Gauge Theories," Phys. Rev. D79: 025002, 2009, arXiv:0807.0163 [hep-th]

[17] J. Figueroa-O'Farrill, G. Papadopoulos, "Pluecker-type relations for orthogonal planes," J. Geom. Phys. 49 (2004) 294, math/0211170]

[18] G. Papadopoulos, "M2-branes, 3-Lie Algebras and Plucker relations," JHEP 0805 (2008) 054, arXiv:0804.2662 [hep-th]

[19] J. P. Gauntlett, J. B. Gutowski, "Constraining Maximally Supersymmetric Membrane Actions," JHEP 0806 (2008) 053, arXiv:0804.3078 [hep-th]

[20] D. Gaiotto, E. Witten, "Janus Configurations, Chern-Simons Couplings, And The Theta-Angle in N=4 Super Yang-Mills Theory," arXiv:0804.2907[hep-th]

[21] Y. Honma, S. Iso, Y. Sumitomo, S. Zhang, "Janus field theories from multiple M2 branes," Phys.Rev. D78 :025027,2008, arXiv:0805.1895 [hep-th].

[22] K. Hosomichi, K-M. Lee, S. Lee, S. Lee, J. Park, "N=5,6 Superconformal Chern-Simons Theories and M2-branes on Orbifolds," JHEP 0809 (2008) 002, arXiv:0806.4977[hep-th]

[23] M. Schnabl, Y. Tachikawa, "Classification of N=6 superconformal theories of ABJM type," arXiv:0807.1102[hep-th]

[24] S. Mukhi, C. Papageorgakis, M2 to D2, JHEP 0805 (2008) 085.

[25] J. Gomis, G. Milanesi, J. G. Russo, "Bagger-Lambert Theory for General Lie Algebras," JHEP 0806: 075, 2008, arXiv:0805.1012 [hep-th].

[26] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni, H. Verlinde, "N=8 superconformal gauge theories and M2 branes," JHEP 0901: 078, 2009, arXiv:0805.1087 [hep-th].

[27] P.-M. Ho, Y. Imamura, Y. Matsuo, "M2 to D2 revisited," JHEP 0807: 003, 2008, arXiv:0805.1202 [hep-th].

[28] M. A. Bandres, A. E. Lipstein, J. H. Schwarz, "Ghost-Free Superconformal Action for Multiple M2-Branes," JHEP 0807: 117, 2008, arXiv:0806.0054 [hep-th]

[29] P. de Medeiros, J. Figueroa-O'Farrill, E. Me'ndez-Escobar, P. Ritter, "On the Lie-algebraic origin of metric 3-algebras," Commun. Math. Phys. 290 (2009) 871, arXiv:0809.1086 [hep-th]

[30] S. A. Cherkis, V. Dotsenko, C. Saeman, "On Superspace Actions for Multiple M2-Branes, Metric 3-Algebras and their Classification," Phys. Rev. D79 (2009) 086002, arXiv:0812.3127 [hep-th]

[31] P.-M. Ho, Y. Matsuo, S. Shiba, "Lorentzian Lie (3-)algebra and toroidal compactification of M/string theory," arXiv:0901.2003 [hep-th]

[32] P. de Medeiros, J. Figueroa-O'Farrill, E. Mendez-Escobar, P. Ritter, "Metric 3-Lie algebras for unitary Bagger-Lambert theories," JHEP 0904: 037, 2009, arXiv:0902.4674 [hep-th]

[33] H. Nishino, S. Rajpoot, Triality and Bagger-Lambert Theory, Phys. Lett. B671 (2009) 415, arXiv:0901.1173.

[34] A. Gustavsson, S-J. Rey, Enhanced N=8 Supersymmetry of ABJM Theory on R(8) and R(8)/Z(2), arXiv:0906.3568 [hep-th].

[35] O. Aharony, O. Bergman, D. L. Jafferis, Fractional M2-branes, JHEP 0811 (2008) 043, arXiv:0807.4924.

[36] M. Hanada, L. Mannelli, Y. Matsuo, Large-N reduced models of supersymmetric quiver, Chern-Simons gauge theories and ABJM, arXiv:0907.4937 [hep-th].

[37] G. Ishiki, S. Shimasaki, A. Tsuchiya, Large N reduction for Chern-Simons theory on S^3, Phys. Rev. D80 (2009) 086004, arXiv:0908.1711.

[38] H. Kawai, S. Shimasaki, A. Tsuchiya, Large N reduction on group manifolds, arXiv:0912.1456 [hep-th].

[39] G. Ishiki, S. Shimasaki, A. Tsuchiya, A Novel Large-N Reduction on S^3: Demonstration in Chern-Simons Theory, arXiv:1001.4917 [hep-th].

[40] Y. Pang, T. Wang, From N M2's to N D2's, Phys. Rev. D78 (2008) 125007, arXiv:0807.1444.

[41] J. DeBellis, C. Saemann, R. J. Szabo, Quantized Nambu-Poisson Manifolds in a 3-Lie Algebra Reduced Model, JHEP 1104 (2011) 075, arXiv:1012.2236.

[42] M. M. Sheikh-Jabbari, Tiny Graviton Matrix Theory: DLCQ of IIB Plane-Wave String Theory, A Conjecture , JHEP 0409 (2004) 017, arXiv:hep-th/0406214.

[43] J. Gomis, A. J. Salim, F. Passerini, Matrix Theory of Type IIB Plane Wave from Membranes, JHEP 0808 (2008) 002, arXiv:0804.2186.

[44] K. Hosomichi, K. Lee, S. Lee, Mass-Deformed Bagger-Lambert Theory and its BPS Objects, Phys.Rev. D78 (2008) 066015, arXiv:0804.2519.

[45] L. Smolin, "M theory as a matrix extension of Chern-Simons theory," Nucl.Phys. B591 (2000) 227, hep-th/0002009.

[46] L. Smolin, "The cubic matrix model and a duality between strings and loops," hep-th/0006137.

[47] T. Azuma, S. Iso, H. Kawai, Y. Ohwashi, "Supermatrix Models," Nucl.Phys. B610 (2001) 251, hep-th/0102168.

[48] J. Palmkvist, "Three-algebras, triple systems and 3-graded Lie superalgebras," J.Phys. A43 (2010) 015205, arXiv:0905.2468 [hep-th]

[49] S. Kim, J. Palmkvist, "N=5 three-algebras and 5-graded Lie superalgebras," J.Math.Phys. 52 (2011) 083502, arXiv:1010.1457 [hep-th]

[50] J. Palmkvist, "Unifying N = 5 and N = 6," JHEP 1105 (2011) 088, arXiv:1103.4860.

[51] B.E.W. Nilsson, J. Palmkvist, "Superconformal M2-branes and generalized Jordan triple systems," Class.Quant.Grav. 26 (2009) 075007, arXiv:0807.5134 [hep-th]

[52] S. P. Chowdhury, S. Mukhopadhyay, K. Ray, "BLG theory with generalized Jordan triple systems," arXiv:0903.2951 [hep-th]

[53] M. Sato, " Covariant Formulation of M-Theory," Int. J. Mod. Phys. A24 (2009), 5019, arXiv:0902.1333 [hep-th]

[54] M. Sato, "Model of M-theory with Eleven Matrices," JHEP 1007 (2010) 026, arXiv:1003.4694 [hep-th]

[55] M. Sato, "Supersymmetry and the Discrete Light-Cone Quantization Limit of the Lie 3-algebra Model of M-theory," Phys. Rev. D85 (2012) 046003, arXiv:1110.2969 [hep-th]

[56] M. Sato, "Zariski Quantization as Second Quantization," Phys. Rev. D85 (2012) 126012, arXiv:1202.1466 [hep-th]

[57] M. Sato, "3-Algebras in String Theory," Linear Algebra - Theorems and Applications, Edited by Hassan Abid Yasser, InTech, Croatia, 2012

[58] M. Sato, "Three-Algebra BFSS Matrix Theory," Int.J.Mod.Phys. A28 (2013) 1350155, arXiv:1304.4430 [hep-th]

[59] M. Sato, "Extension of IIB Matrix Model by Three-Algebra," Int.J.Mod.Phys. A28 (2013) 1350083, arXiv:1304.4796 [hep-th]

[60] M. Sato, "Four-algebraic extension of the IIB matrix model," PTEP 2013 (2013) 7, 073 B04, arXiv:1304.7904 [hep-th]

[61] N.Kamiya,;A structure theory of Freudenthal-Kantor triple systems,J.Alg. 110 (1987) 108-123.

[62] N.Kamiya,;A structure theory of Freudenthal-Kantor triple systems II,Comm. Math.Univ.Sancti Pauli.,38(1989)41-60.

[63] N.Kamiya,;A structure theory of Freudenthal-Kantor triple systems III, Mem.Fac. Sci.Shimane Univ. 23(1989)33-51.

[64] N.Kamiya,;The construction of all Simple Lie algebras over C from balanced Freudenthal-Kantor triple systems, Contributions to General Algebra 7,Verlag Hoder-Pichler-Tempsky, Wien,Veralg.G.Teubner,Stutugart(1991) 205-213.

[65] N.Kamiya,;On Freudenthal-Kantor triple systems and generalized structurable algebras, Proceeding in International conference of nonassociative algebras and its applications, Mathematics and Its Applications 303(1994) 198-203,Kluwer Academic Publisher,

[66] N.Kamiya,;On the Peirce decompostions for Freudenthal-Kantor triple systerns, Comm. in Alg. 25,(6),(1997),1833-1844.

[67] N.Kamiya, Examples of Peirce decomposition of generalized Jordan triple system of second order,-balanced cases-, Contemmporary Math., vol.391,AMS.(2005), Noncommutative Geometry and Representation theory an Mathematical Physics,(ed)Fuchs, p157-165.

[68] N.Kamiya and S.Okubo, On δ-Lie supertriple systems associated with (ε, δ)-Freudenthal-Kantor triple systems, Proc. Edinb. Math.Soc. 43(2000),243-260.

[69] I.L.Kantor and N.Kamiya, A Peirce decomposition for generalized Jordan triple systems of second order, Comm in Alg.,31(2003),5875-5913.

[70] I.L.Kantor,;Models of exceptional Lie algebras, Soviert Math.Dokl.14(1973) 254-258.

[71] I.L.Kantor,;Some generalization of Jordan algebras,Trudy Sem.Vektor, Tensor Ana1.16(1972)407-499(Russian).

[72] W.Kaup,;Algebraic characterization of symmetric complex Banach manifoleds, Math. Ann.228. (1977)39-64.

[73] I.Loos,;Bounded symmetric domains and Jordan pairs,Mathematical Lectures.lnvine: University of California at Irwine,1977.

[74] K.Meyberg,;Lecture on algebras and triple systems,Lecture Notes,the Univ. ofVirginia,1972.

[75] E.Neher,;On the radical of J*-triples, Math.Z.211. (1992)323-332.

[76] B.N.Allison,; A class of nonassociative algebras with involution containing class of Jordan algebras, Math.Ann., 1978,237(2),133-156.

[77] N.Jacobson,;Structure and representations of Jordan algebras. Amer. Math. Soc.,Providence.R.1.,1968.

[78] S.Okubo,;lntroduction to Octonion and Other Non-associative Algebras in Physics, Cambridge Univ.Press.1995.

[79] 1.Satake,;Algebraic structure of symmetric domains,Princeton University press.1980.

[80] R.D.Shafer,;An introduction to Non-associative Algebras,Academicpress, 1966.

[81] N.Kamiya and D.Mondoc; A new class of nonassociative algebras with involution, Proc.Japan Acad.,84 Ser.A(2008),no5,68-72,

[82] N.Kamiya and S.Okubo,; A construction of simple Lie superalgebras of certain type from triple systems, Bull.Australia Math.Soc. vol.69,(2004) 113-123.

[83] N.Kamiya and S.Okubo,; Construction of Lie superalgebrasD(2,1,α), $G(3)$ and $F(4)$ from some triple systems. Proc. Edinburgh Math. Soc. (2003) 46, 87-98.